



## Antimicrobial activity of honey samples from different sites in Northeast Algeria

Neila Nedji<sup>1</sup>, Wahida Loucif-Ayad<sup>1,2</sup>

<sup>1</sup>: Laboratory of Applied Animal Biology, Department de Biology, Faculty of Sciences, Badji Mokhtar University, 23000, Annaba, Algeria.

<sup>2</sup>: Faculty of Medicine, Badji Mokhtar University, 23000-Annaba, Algeria.  
[wahloucif@yahoo.fr](mailto:wahloucif@yahoo.fr)

**Abstract:** The purpose of the study was to characterize the physicochemical properties and the antibacterial activity of honey samples collected from different sites of Northeast Algeria. The antibacterial activity of honey against *Bacillus cereus* (IPA), *Staphylococcus aureus* (ATCC25923R), *Escherichia coli* (ATCC25922) and *Pseudomonas aeruginosa* (ATCC27893R) was evaluated by the disc diffusion method and determined as an equivalent of the inhibition zones diameters after incubation of the cultures at 37°C for 24 hours. The moisture content, pH, Electrical conductivity and Ash were measured and the investigation of the polyphenols and flavonoid contents were done spectrophotometrically in each honey sample. Results showed that Algerian honey inhibited the growth of all examined microorganisms with the highest antimicrobial activity against the Gram positive bacteria. Physicochemical parameters were similar between the honey samples collected from different sites and polyphenol and flavonoids contents were variable, depending on the honey samples. The strong antimicrobial activity of Algerian honey may be due to high total phenolic and flavonoid contents and this study suggests potential use of honey in foods.

[Nedji N, Loucif-Ayad W. **Antimicrobial activity of honey samples from different sites in Northeast Algeria.** *Life Sci J* 2021;18(1):73-79]. ISSN: 1097-8135 (Print) / ISSN: 2372-613X (Online). <http://www.lifesciencesite.com>. 10. doi:[10.7537/marslsj180121.10](https://doi.org/10.7537/marslsj180121.10).

**Keywords:** Algerian honey; antimicrobial activity; polyphenols and flavonoid contents

### 1. Introduction

Microbial infections are the cause of a large burden of diseases and bacteria are listed in the first position among the common microorganisms responsible for opportunistic diseases. Therapy of bacterial infections is a frequent problem due to the emergence of bacterial strains resistant to numerous antibiotics [1]. Food-borne diseases caused by the consumption of contaminated foods have a wide economic and public health impact worldwide. Many pathogenic microorganisms (*Staphylococcus aureus*, *Bacillus cereus*, *Escherichia coli*, and *Pseudomonas aeruginosa*) have been reported as the causal agents of food-borne diseases [2, 3]. A variety of different chemical and synthetic compounds have been used as antimicrobial agents to inhibit bacteria in foods. There is great interest in controlling the growth or eliminating foodborne pathogens using natural antimicrobials [4].

Honey, a natural product formed from nectar by honeybees, has attracted much attention in recent years as an useful ingredient applied in medicine [5-7]. Honey possesses biological properties including antioxidant [8], fungicidal [9], anti-inflammatory [10], reproductive [11-12], hypoglycemic [8] and

antibacterial effects [13]. Numerous reports describe the antibacterial properties of honey but there has been only limited research on antimicrobial activity of Algerian honey [14-16].

Antimicrobial activity of honey varies greatly with processing and origin, depending on the natural vegetative flowers blooming in different seasons and in different places [13, 17]. The healing effect of honey could be due to various physical and chemical properties [18]. The floral source of honey plays an important role on its biological properties [19, 20]. An extensive review of the antimicrobial activity of honey showed it to be derived from high sugar content low water content, acidity, the generation of hydrogen peroxide on dilution [21-23] and phytochemical components [18,24], aromatic acids and phenolic compounds [25].

The present investigation was undertaken to evaluate the antibacterial potential of Algerian honey against a range of pathogenic bacteria and its physicochemical properties.

## 2. Material and Methods

### Honey samples and preparation

Honey samples were gathered from honeybee colonies of the local strain *Apis mellifera intermissa* in four sites of Annaba, in Northeast Algeria: Seraidi (36°54'39.92"N, 7°37'36.24"E), Chetaibi (36°59'41.47"N, 7°19'30.29"E), Berrehal (36°50'16.89"N, 7°26'56.51"E) and El-Bouni (36°49'35.92"N, 7°39'51.45"E), and designed as SH, CH, BH and EH respectively. The four sampling sites belong to the same climatic level (sub-humide). Seraidi and Chetaibi are located far from road traffic and presented a rich floral diversity as compared to Berrehal and El-Bouni. All the samples were collected in september 2013 from the nests using sterile syringes, stored in dark glass bottles at room temperature. Initially, the honeys were subjected to sterilization by  $\gamma$ -irradiation at a dose of 25 kGy [26].

### Physicochemical analyses

The determination of moisture [27] was ascertained by refractometry. All measurements were performed at 20 °C, after waiting for 6 min for equilibrium, and obtaining the corresponding % moisture from the refractive index of the honey sample by consulting a standard table for the purpose [28].

The pH value was determined in a solution containing 10 g of honey in 75 mL of distilled water. Determination of pH was done with pH-meter mode [27].

Honey Electrical conductivity was determined by conductimetric assay from a solution containing 10 g of honey in 75 ml of distilled water [29].

Ash content was indirectly determined using the measured Electrical conductivity and applying the following equation:  $X1 = (X2 - 0.143) / 1.743$ , where:  $X1$  = ash value;  $X2$  = Electrical conductivity in mS/cm at 20 °C [30].

All physicochemical tests were performed in triplicate.

### Polyphenols of honey

Total polyphenol contents of honey were determined by the Folin–Ciocalteu colorimetric method [31–32]. Honey sample (0.1 mL) was mixed with 0.5 mL of the Folin–Ciocalteu reagent and 0.5 mL of 7.5% (w/v)  $\text{Na}_2\text{CO}_3$ , and the reaction was kept

in the dark for 1h, after which the absorbance was read at 725 nm. The total polyphenol content was calculated based on a standard curve prepared using gallic acid and expressed as milligrams of gallic acid equivalent (GAE) per 100 gram of sample.

### Flavonoids of EEP

Total flavonoid contents in honey were determined by the method of [33] with minor modifications [32]. To 0.1 mL of the honey sample solution (10% w/v), 0.5 mL of 20 mg  $\text{AlCl}_3/\text{mL}$  ethanol 96% (v/v) solution was added. After 1h at room temperature, the absorbance was measured at 420 nm. Total flavonoids are calculated as mg quercetin equivalents QE/100 g honey from a calibration curve.

### Antimicrobial activity test

Antimicrobial activity of honey samples were investigated by the disc diffusion method [34]. The antimicrobial screening was performed using Mueller-Hinton agar. The bacteria tested were graciously provided by Pasteur Institute of Algiers (Algeria) and included two Gram-positive bacteria [*Bacillus cereus* (IPA): *Staphylococcus aureus* (ATCC 25923R)] and two Gram-negative bacteria [*Escherichia coli* (ATCC25922), *Pseudomonas aeruginosa* (ATCC 27893R)] strains. Agar disc diffusion method was employed for the determination of antimicrobial activities of Algerian honey. Suspensions of tested microorganisms (0.5 Mac Farland scale) were spread into solid media plates. Filter paper discs (6mm in diameter) were impregnated with 20 $\mu$ l of each honey sample and with distilled water (control) and the inoculated plates were incubated at 37°C for 24 hours. Diameters of the inhibition zones were measured in millimeters. All the tests were performed in triplicate.

### Statistical analysis

The results are reported as mean  $\pm$  standard deviation (m  $\pm$  SD). One-way ANOVA and Tukey post hoc multiple comparison tests were used to analyze data. P value less than 0.05 were considered significant.

## 3. Results

The results of the analysis of quality parameters, such as moisture, pH, Electrical conductivity and Ash content are summarized in Table 1.

Table1. Collection sites and physicochemical parameters of honey samples. Data are expressed as means  $\pm$  SD.

Honey	Collection site	Moisture content (%)	pH	Electrical conductivity (mS/cm)	Ash (%)
SH	Seraidi	17.88 $\pm$ 0.22 <sup>a</sup>	3.77 $\pm$ 0.05 <sup>a</sup>	0.36 $\pm$ 0.06 <sup>a</sup>	0.12 $\pm$ 0.03 <sup>a</sup>
CH	Chetaibi	17.94 $\pm$ 0.63 <sup>a</sup>	3.83 $\pm$ 0.05 <sup>a</sup>	0.29 $\pm$ 0.05 <sup>a</sup>	0.08 $\pm$ 0.03 <sup>a</sup>
BH	Berrehal	18.24 $\pm$ .60 <sup>a</sup>	3.89 $\pm$ 0.18 <sup>a</sup>	0.38 $\pm$ 0.11 <sup>a</sup>	0.14 $\pm$ 0.06 <sup>a</sup>
EH	El-Bouni	18.38 $\pm$ 0.51 <sup>a</sup>	4.6 $\pm$ 0.08 <sup>a</sup>	0.41 $\pm$ 0.02 <sup>a</sup>	0.15 $\pm$ 0.01 <sup>a</sup>

Means with the same superscript letters within a column are not significantly different (P > 0.05).

There were no significant differences ( $P > 0.05$ ) in all physicochemical parameters between honey samples of different sites. Moisture is a parameter related to the maturity degree of honey and temperature. In the present study, moisture values are between 17.88% and 18.38% and all the Algerian honeys analyzed were found to be acidic in character. Their pH values ranged from 3.77 to 4.6 (Table 1).

The Electrical conductivity (mS/cm) in honey samples, varied in the range of 0.29 to 0.41 and the Ash content is generally small and its values were ranged from 0.08 to 0.15 %.

#### Total polyphenols and flavonoid contents

Total polyphenol and flavonoid contents of honey samples from the four sites of Algeria were investigated (Table 2).

**Table 2.** Total polyphenols and flavonoids contents in honey samples.

Honey	Total polyphenol* (mg GAE/100g of honey)	Total flavonoid** (mg EQ /100g of honey)
SH	189.2 ± 16.01 <sup>a</sup>	95.94 ± 5.97 <sup>a</sup>
CH	131.3 ± 13.64 <sup>b</sup>	69.99 ± 7.36 <sup>b</sup>
BH	99.03 ± 3.97 <sup>b c</sup>	64.47 ± 1.39 <sup>b c</sup>
EH	92.14 ± 5.23 <sup>c</sup>	44.56 ± 3.88 <sup>c</sup>

\*Total polyphenol contents were determined by the Folin–Ciocaltau method. Value is mean ± standard deviation.

\*\*Flavonoid contents were determined by  $AlCl_3$  coloration.


Means with different superscript letters within a column are significantly different at  $P < 0.05$  (ANOVA followed by a post-hoc Tukey test).

Results showed that there was a significant difference between the sites for total polyphenol contents ( $F=16.48$ ,  $df= 3$ ,  $P<0.001$ ) and also for flavonoid contents ( $F=15.42$ ,  $df= 3$ ,  $P<0.001$ ).

The total polyphenol contents of honey studied were ranged between 92.14 to 189.2 mg GAE/100g of honey. Honeys also contained flavonoids at levels of 44.56-95.94 mgQE/100g of honey, with the higher values observed in honey samples from Seraidi and Chetaibi sites.

#### Antibacterial Activity Assay

The disc diffusion method was used to determine the inhibition zones of the different honey from the four sites. The two Gram positive and two Gram negative bacteria have been used. According to the results in figure1, different honey samples showed antibacterial activity against all bacteria with the highest antibacterial activity against the Gram positive bacteria.



**Figure 1.** Diameters of the zones of inhibition (mm) of the growth bacteria tested according to the honey samples.

In our results, the antimicrobial activity of honey from the four sites varies according to the origin of the honey. The largest inhibitory zones of the growth bacteria were noticed for the honey of Seraidi and Chetaibi sites which showed also the highest values of polyphenol and flavonoid contents.

#### 4. Discussions

In the present study, the analysis of some parameters of Algerian honey, such as moisture, pH, Electrical conductivity and Ash content was evaluated. Moisture values were within the values found in Moroccan honeys by [35] (14.64% - 18.59%) and [36] (14.5% - 23.6%). The moisture content of honey depends on various factors such as harvesting season, degree of maturity reached in the hive and climatic factors [37]. Also, the moisture content of honey is a highly important factor contributing to its stability against fermentation and granulation during storage [38, 39].

In general, honey is acidic in nature, irrespective of its variable geographical origin. The pH values of Indian, Morocco, Argentinean honeys and Saudi honeys, have been found to vary between 3.7 to 4.4, 3.91 to 4.93, 3.25 to 3.32 and 3.48 to 6.06, respectively [40,41]. This parameter is of great importance, during extraction and storage of honey, as it influences the texture, stability and shelf life of honey [42]. The values obtained in this study were similar to those previously reported for other honey samples from India, Brazil, Spain and Turkey, which were reported to have pHs between 3.49 and 4.70 [39,40]. The pH values of Algerian honey samples previously reported were ranged from 3.49 to 4.43 [41], 3.29 to 4.37 [42] and 3.70 to 4.00 [43].

A linear relationship is known to exist between the electrical conductivity and the Ash content. Similarly, a correlation has been found between the electrical conductivity and Ash content for some Algerian honeys [41, 43]. The electrical conductivity shows great variability according to the floral origin, and is important for differentiating honeys of different floral origin [44]. Ash content is a parameter that has been associated with botanical and geographical origins of honey samples. The Ash content in honey is generally small and depends on nectar composition of predominant plants in their formation [45]. Our results are in agreement with those of some authors [46, 47]. Polyphenols are an important group of compounds which influence the appearance and the functional properties of honey [48, 49].

The total polyphenolic and flavonoid content of honey samples has been determined in different countries. [50] has reported that total phenols in Chinese honey were 9.41 to 102.1 mg GAE/100 g and in Romanian honey samples, values were 23.0-125.0 mg GAE/100 g [38,51]. Total phenols of Burkina Faso honey were 32.59-114.75 mg GAE/100 g [52] while in Paraguay, it varied between 125.17 and 176.50 mg GAE/100g honey [32]. For Indian and Croatian honeys, the phenolic content ranged from 48 to 99 and 31.72 to 80.11 mg GAE/ 100 g, respectively [53,39]. Total phenols of Algeria were 63.93 - 95.36 mg GAE /100g [1].

The concentration and type of polyphenolic substances depend on the floral origin of honey and are major factors responsible for biological activities [54, 55], including antimicrobial activities [56]. Also, flavonoid contents in Romanian honey samples were 38.96-65, 98 mg QE/100 g of honey [54]. In samples from Venezuela, flavonoid content were ranged from 2.6 to 31.0 mg quercetin equivalents/100g of honey [57] and [58] has reported that total flavonoid content ranged from 6.9 to 28.5 mg quercetin equivalents per 100 g. Total flavonoid contents varied from 41.88 to 211.68 mg quercetin equivalent kg<sup>-1</sup> of Italian honeys [59] and flavonoids varied between 2.52 – 27.21mgQE/100g for honey samples of Nigeria. Flavonoid contents were ranged between 27.07-71.78 mg /kg of honey in sample from Algeria [43].

In this study, the antimicrobial activity of honey was investigated and showed antibacterial activity against all bacteria tested with high antimicrobial activity against Gram+ bacteria. The antimicrobial activity of the honey from the four sites varies according to the origin of the honey. The honey samples of Seraidi and Chetaibi had strong antimicrobial activity as compared to those from Berrahel and El-Bouni. This strong antimicrobial activity is related with high values of total polyphenol and flavonoid contents in SH and CH. The rich floral

diversity of the two sites (Seraidi and Chetaibi) seems to play an important role in polyphenol and flavonoid contents, influencing the honey's antimicrobial activity. In fact, honey from different regions varied in its ability to inhibit the growth of bacteria suggesting that botanical origin plays an important role in influencing a honey's antimicrobial activity [4,25,60,61]. The total content of phenolic compounds is dependent on the botanical source of honey and its collection region [58].

Natural products are promising natural antimicrobial agents with potential applications in pharmaceutical or food industries for controlling the pathogenic bacteria. The strong antibacterial effects of Algerian honey against foodborne pathogens such as *B. cereus* and *S. aureus* suggests potential use as a food preservative against pathogenic food-related microorganisms. Other research will be pursued to determine the plant origin of Algerian honey and its qualitative chemical composition.

#### Acknowledgements:

Foundation item: CNEPRU project to Dr. W. Loucif-Ayad). [grant No.F011 2011 / 00097]. Authors are grateful to the Ministry of High Education and Scientific Research of Algeria for financial support to carry out this work.

#### Corresponding Author:

Dr. Wahida Loucif-Ayad  
Badji Mokhtar University, 23000, Annaba, Algeria  
E-mail: [wahloucif@yahoo.fr](mailto:wahloucif@yahoo.fr)

#### References

1. Ahmed M, Djebli N, Aissat S, Zerrouki K & Bourabeh A. In Vitro synergistic antibacterial activity of natural honey combined with curcuma starch and their correlation with diastase number, flavonoid and polyphenol content. J Plant Pathol Microb 2013; 4 152. doi:10.4172/2157-7471.1000152
2. Erkmen O. Antimicrobial effects of Turkish propolis, pollen, and laurel on spoilage and pathogenic food-related microorganisms. J Med Food 2008;7: 587–592.
3. El-Bassiony TA, Saad NM, El-Zamkan MA. Study on the antimicrobial activity of Ethanol Extract of Propolis against Enterotoxigenic Methicillin-Resistant *Staphylococcus aureus* in lab prepared Ice-cream. Vet World 2012;5 (3): 155-159.
4. Taormina PJ, Niemira B.A., Beuchat LR. Inhibitory activity of honey against foodborne pathogens as influenced by the presence of hydrogen peroxide and level of antioxidant



- power. *Int J Food Microbiol* 2001; 69(3): 217-225.
5. Mandal S, DebMandal M, Pal NK, Saha K. Antibacterial activity of honey against clinical isolates of *Escherichia coli*, *Pseudomonas aeruginosa* and *Salmonella enterica* serovar *Typhi*. *Asian Pac J Trop Med* 2010; 3(12): 961-964.
  6. DebMandal M, Mandal S. Honey: its medicinal property and antibacterial activity. *Asian Pac J Trop Biomed* 2011; 1(2): 154-160.
  7. Alvarez-Suarez JM, Giampieri F, Battino M. Honey as a source of dietary antioxidants: structures, bioavailability and evidence of protective effects against human chronic diseases. *Curr Med Chem* 2013; 20(5): 621-638.
  8. Erejuwa OO, Gurtu S, Sulaiman SA, AbWahab MS, Sirajudeen KN, Salleh MS. Hypoglycemic and antioxidant effects of honey supplementation in streptozotocin-induced diabetic rats. *Int J Vitam Nutr Res* 2010; 80: 74-82.
  9. Koc AN, Silici S, Kasap F, Hormet-Oz HT, Mavus-Buldu H, Ercal BD. Antifungal activity of the honeybee products against *Candida spp.* and *Trichosporon spp.* *J Med Food* 2011; 14: 128-134.
  10. Kassim M, Achoui M, Mustafa MR, Mohd MA, Yusoff KM. Ellagic acid, phenolic acids and flavonoids in Malaysian honey extracts demonstrate *in vitro* anti-inflammatory activity. *Nutr Res* 2010; 30: 650-659.
  11. Mohamed M, Sulaiman SA, Jaafar H, Sirajudeen KN. Effect of different doses of Malaysian honey on reproductive parameters in adult male rats. *Andrologia* 2011; doi:10.1111/j.1439-0272.2010.01159.x.
  12. Zaid SS, Sulaiman SA, Sirajudeen KN, Othman NH. The effects of Tualang honey on female reproductive organs, tibia bone and hormonal profile in ovariectomised rats-animal model for menopause. *BMC Complement Altern Med* 2011;10: 82.
  13. Tan HT, Rahman RA, Gan SH, Halim AS, Hassan SA, Sulaiman SA, Kirnpal-Kaur B. The antibacterial properties of Malaysian tualang honey against wound and enteric microorganisms in comparison to manuka honey. *BMC Complement Altern Med* 2009; 9: 34.
  14. Ouchemoukh S, Schweizer P, Bachir Bey M, Djoudad-Kadji H, Louaileche H. HPLC sugar profiles of Algerian honeys. *Food Chem* 2010; 121: 561-568.
  15. Ahmed M, Aissat S, Djebli N, Boulkaboul A, Meslem AM, et al. The Influence of Starch of Ginger on the Antibacterial Activity of Honey of Different Types from Algeria against *Escherichia coli* and *Staphylococcus aureus*. *IJMR* 2011, 2: 258-262.
  16. Ahmed M, Djebli N, Aissat S, Meslem AM, Bacha S. The Influence of botanical origin and physicochemical parameters on the antifungal activity of Algerian honey. *J Plant Pathol Microb* 2012; 3: 5.
  17. Manyi-Loh CE, Clarke AM, Ndip RN. An overview of honey: therapeutic properties and contribution in nutrition and human health. *Afr J Microbiol Res* 2011; 5: 844-852.
  18. Snow M, Manley-Harris M. On the nature of non-peroxide antibacterial activity in New Zealand Manuka honey. *Food Chem* 2004; 84: 145-147.
  19. Bogdanov S. Harmonized Methods of the European Honey Commission. International Honey Commission 2002.
  20. Molan, PC. Not all honeys are the same for wound healing. *Bull. Eur. Tissue Rep Soc* 2002; 9: 5-6.
  21. Kačániová M, Fatrcova-Sramkova K, Nozkova J, Melich M, Kadasi-Horakova M, Knazovicka V, Felsociova S, Kunova S, Mariassyova M. Antiradical activity of natural honeys and antifungal effect against *Penicillium genera*. *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 2011; 46 (1): 92-96.
  22. Hegazi AG, Abd Allah FM. Antimicrobial activity of different Saudi Arabia honeys. *Global Veterinaria* 2012; 9 (1): 53-59.
  23. Mistry R, Shah G. Study of inhibitory effect of honey on various pathogenic and non-pathogenic microorganisms *Int. J Appl Sci Biotechnol* 2013; 1(4): 279-281.
  24. Kwakman PH, Zaat SA. Antibacterial Components of Honey. *IUBMB Life* 2012; 64: 48-55.
  25. Alvarez-Suarez JM, Tulipani S, Díaz D, Estevez Y, Romandini S, Giampieri F, Damiani E, Astolfi P, Bompadre S, Battino M. Antioxidant and antimicrobial capacity of several monofloral Cuban honeys and their correlation with color, polyphenol content and other chemical compounds. *Food Chem Toxicol* 2010; 48(8-9): 2490-2499.
  26. Molan PC, Allen KL. The effect of gamma-irradiation on the antibacterial activity of honey *J Pharm Pharmacol* 1996; 48(11): 1206-1209.
  27. AOAC. Official Methods of Analysis, 15th ed. Association of Official Analytical Chemists, 1990; Inc., Arlington.
  28. Zerrouk SH, Fallico BG, Arena EN, Ballistreri GF, boughediri LA. Quality evaluation of some honey from the central region of Algeria. *Jordan*

- Journal of Biological Sciences 2011; 4(4): 243-248.
29. Sancho MT, Muniategui S, Huidobro JF, Simal J. Evaluating soluble and insoluble ash, alkalinity of soluble and insoluble ash and total alkalinity of Ash in honey using electrical conductivity measurements at 20°C. *Apidologie* 1992; 23: 291–297.
  30. Piazza MG, Accorti M, Persano Oddo L. Electrical conductivity, ash, colour and specific rotatory power in Italian unifloral honeys. *Apicoltura* 1991;7: 51–63.
  31. Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin–Ciocalteu reagent. *Methods of Enzymology* 1999; 299: 152-178.
  32. Vit P, Gutiérrez MG, Rodríguez-Malaver AJ, Aguilera G, Fernández-Díaz C, Tricio AE. Comparación de mieles producidas por la abeja yateí (*Tetragonisca fiebrigi*) en Argentina y Paraguay. *Acta Bioquímica Clínica Latinoamericana* 2009; 43: 219-226.
  33. Woisky RG, Salatino A. Analysis of propolis: some parameters and procedures for chemical quality control. *Journal of Apicultural Research* 1998; 37: 99-105.
  34. Bauer AW, Kirby MDK, Sherris JC, Truck M. Antibiotic susceptibilities testing by standard single disc diffusion method. *American Journal of Clinical Pathology* 1966;45: 493-496.
  35. Chakir A, Abderrahmane R, Gian LM, Paola F. Physicochemical properties of some honeys produced from different plants in Morocco. *Arabian Journal of Chemistry* 2011; doi:10.1016/j.arabjc.2011.10.013.
  36. Terrab A, Gustavo González A, Díez MJ, Heredia FJ. Characterisation of Moroccan unifloral honeys using multivariate analysis. *Eur Food Res Technol* 2003; 218: 88-95.
  37. Finola MS, Lasagno MC, Marioli JM. Microbiological and chemical characterization of honeys from central Argentina. *Food Chemistry* 2007; 100: 1649- 1653.
  38. Al ML, Daniel D, Moise A, Bobis O, Laslo L, Bogdanov S. Physico-chemical and bioactive properties of different floral origin honeys from Romania. *Food Chemistry* 2009; 112: 863–867.
  39. Saxena S, Gautam S, Sharma A. Physical, biochemical and antioxidant properties of some Indian honeys. *Food Chemistry* 2010; 118: 391–397.
  40. Azeredo LDC, Azeredo M, De Souza SR, Dutra VML. Protein content and physicochemical properties in honey samples of *Apis mellifera* of different floral origins. *Food Chemistry* 2003; 80: 249-254.
  41. Ouchemoukh S, Louaileche H, Schweizer P. Physicochemical characteristics and pollen spectrum of some Algerian honey. *Food Control* 2007;18: 52-58.
  42. Chefrour C, Draiaia R, Tahar A, Ait Kaki Y, Bennadja S, Battesti M. Physicochemical characteristics and pollen spectrum of some north-east Algerian honeys. *Afr J Food Agric Nutr Dev* 2009; 9: 1276-1293.
  43. Khalil MI, Moniruzzaman M, Boukraâ L, Benhanifia M, Islam MA, Islam MN, Sulaiman SA, Gan SH. Physicochemical and Antioxidant Properties of Algerian Honey. *Molecules* 2012; 17: 11199-11215.
  44. Terrab A, Recamales AF, Hernanz D, Heredia FJ. Characterisation of Spanish thyme honeys by their physicochemical characteristics and mineral contents. *Food Chem* 2004; 88: 537-542.
  45. Adams CJ, Boulton CH, Deadman BJ, Farr JM, Grainger MN, et al. Isolation by HPLC and characterization of the bioactive fraction of New Zealand manuka (*Leptospermum scoparium*) honey. *Carbohydr Res* 2008; 343: 651-659.
  46. Al-Khalifa AS, Al-Arifia IA. Physicochemical characteristics and pollen spectrum of some Saudi honeys. *Food Chemistry* 1999; 67: 21-25.
  47. Sudhanshu S, Satyendra G, Arun S. Physical, biochemical and antioxidant properties of some Indian honeys. *Food Chem* 2010; 118: 391-397.
  48. Cimpoiou C, Hosu A, Miclaus V, Puscas A. Determination of the floral origin of some Romanian honeys on the basis of physical and biochemical properties. *Spectrochim Acta A Mol. Biomol Spectrosc* 2012; 100: 149-154.
  49. Khalil MI, Alam N, Moniruzzaman M, Sulaiman SA, Gan SH. Phenolic acid composition and antioxidant properties of Malaysian honeys. *J Food Sci* 2011; 76: 921-928.
  50. Dong R, Zheng YN, Xu BJ. Phenolic profiles and antioxidant capacities of Chinese unifloral honeys from different botanical and geographical sources. *Food Bioprocess Tech* 2013; 6(3): 762-770.
  51. Bobis O, Mărghitas LA, Dezmirean D, Chirilă F, Moritz RFA. Preliminary Studies Regarding Antioxidant and Antimicrobial Capacity for Different Types of Romanian Honeys. *Bulletin UASVM Animal Science and Biotechnologies* 2011; 68(1-2): 91-97.
  51. Meda A, Lamien CE, Romito M, Millogo J, Nacoulma OG. Determination of the total phenolic, flavonoid and proline contents in Burkina Fasho honey, as well as their radical

- scavenging activity. *Food Chem* 2005; 91: 571-577.
52. Krpan M, Marković K, Šarić G, Skoko B, Hruškar M, et al. Antioxidant activities and total phenolics of acacia honey. *Czech J Food Sci* 2009; 27: 245-247.
  53. Bobis O, Mărghitas LA, Dezmirean D, Bonta V, Mihai CM. Beehive products: source of nutrients and natural biologically active compounds. *J Agroalim Proc Technol* 2010; 16(2): 104-109.
  54. Mărghitas LA, Dezmirean D, Pocol C, Ilea M, Bobis O, Gergen I. The development of a biochemical profile of acacia honey by identifying biochemical determinants of its quality. *Notulae Botanicae Horti Agro Botanici Cluj-Napoca* 2010; 38(2): 9-13.
  55. Al-Mamarya M, Al-Meerib A, Al-Haborib M. Antioxidant activities and total phenolics of different types of honey. *Nutrition Research* 2002; 22: 1041-1047.
  56. Rodríguez-Malaver AJ, Rasmussen C, Gutiérrez MG, Gil F, Nieves B, Vit P. Properties of honey from ten species of Peruvian stingless bees. *Natural Product communications* 2009; 4(9): 1221-1226.
  57. Socha R, Juszcak L, Pietrzyk S, Fortuna T. Antioxidant activity and phenolic composition of herb honeys. *Food Chem* 2009; 113:568-574.
  58. Pichichero E, Canuti L, Canini A. Characterization of the phenolic and flavonoid fractions and antioxidant power of Italian honeys of different botanical origin. *J Sci Food Agric* 2009; 89: 609-16.
  59. Silici S, Sagdic O, Ekici L. Total phenolic content, antiradical, antioxidant and antimicrobial activities of Rhododendron honeys. *Food Chemistry* 2010; 121: 238-243.
  60. Liu JR, Ye YL, Lin TY, Wang YW, Peng CC. Effect of floral sources on the antioxidant, antimicrobial, and anti-inflammatory activities of honeys in Taiwan. *Food Chemistry* 2013; 139 (1-4): 938- 943.

1/15/2021