



Assessment of Genetic Diversity Among Sunflower Inbred Lines for the Development of Recombinants having Better Achene Yield and Oil Quality

Muhammad Aftab¹, Rizwana Qamar¹, Sajida Habib¹, Momna Mehmood², Muhammad Aslam¹, Ali Hassan¹, Amir Bibi², Shamsa Kanwal³, Hafiz Saad Bin Mustafa^{1*}, Shazia Saeed¹, Ahmad Nawaz Gill¹

¹Oilseeds Research Institute, Faisalabad (Pakistan).

²Department of Plant Breeding and Genetics, University of Agriculture Faisalabad, (Pakistan).

³Agricultural Biotechnology Research Institute, Faisalabad (Pakistan).

*Corresponding Author's Email: saadpbg@gmail.com

ABSTRACT: Assessment of genetic diversity in germplasm is prerequisite for the success of any plant breeding program. Twenty-four Sunflower inbred lines including 15 cytoplasmic male sterile lines (A lines) and nine fertility restorer lines (R lines) were sown to estimate genetic dissimilarities at research area of Oilseeds Research Institute, Faisalabad (Pakistan) during Spring 2018. The objective of study was identification of superior inbred lines to be utilize as parental inbred lines for the development of better recombinants and estimation of association among important qualitative and quantitative characters. Data revealed that Sunflower inbred lines possessed noticeable genetic variation and have potential for utilization in hybrid development program. Among A lines, ORI-92, ORI-94 and ORI-25 has proved their worth for good seed yield and can be employed in crossing with R lines RL-66, RL-68 and RL-50 having high Oleic acid content may give a better recombinant as they displayed high degree of genetic divergence among them. Moreover, yield had positive relationship with head diameter and Linoleic acid while negative relationship with Oleic acid.

[Muhammad Aftab, Rizwana Qamar, Sajida Habib, Momna Mehmood, Muhammad Aslam, Ali Hassan, Amir Bibi, Shamsa Kanwal, Hafiz Saad Bin Mustafa, Shazia Saeed, Ahmad Nawaz Gill. **Assessment of Genetic Diversity Among Sunflower Inbred Lines for the Development of Recombinants having Better Achene Yield and Oil Quality**. *Life Sci J* 2022;19(7):38-43]. ISSN 1097-8135 (print); ISSN 2372-613X (online). <http://www.lifesciencesite.com>. 05. doi:[10.7537/marslsj190722.05](https://doi.org/10.7537/marslsj190722.05).

Keywords: Genetic diversity, Hybrids, Inbred lines, Oil quality, Sunflower, Yield

INTRODUCTION

Edible oil is Pakistan's largest food import commodity ranking third in the import list after petroleum products and machinery. Pakistan is dependent upon other countries to fulfill more than 80% edible oil requirement and spending a big amount of foreign exchange every year. The local production of edible oil is less than 20% of country's requirements (Aftab *et al.*, 2019). Domestically 0.50 million tons of edible oil was produced during the year 2018-19 whereas 2.421 million tons of oil worth Rs. 192.20 billion was imported to undertake the constantly growing demand of edible oil (Economic Survey of Pakistan, 2018-19). This situation is posing huge burden to national economy. Considering the importance of local edible oil production, Government of Pakistan has taken initiative to enhance local production by increasing the area under oilseed crops and availability of high yielding hybrids/varieties of oilseed crops. Sunflower

is an important oilseed crop of Pakistan and ranked 2nd after Rapeseeds and Mustards. Moreover, its oil quality with greater content of unsaturated fatty acids also increase its importance from health point of view in current scenario of increasing cardiac issues. Sunflower has a short growing period with wider adaptability which makes it suitable for different cropping pattern and agro-ecological zones of Pakistan (Aftab *et al.*, 2019).

Sunflower hybrids had more importance in agricultural system of Pakistan because of its higher yield, stability and disease resistance but unfortunately no local sunflower hybrid is cultivated in the field at this time. Farmers had to sow internationally developed hybrids which costs a lot and sometimes not adapted well to the local environmental conditions. For this purpose, locally developed sunflower hybrids are direly needed which is only possible through the utilization of locally developed genetic stock of inbred lines. The genetic

stock of inbred lines (A, B and R lines) having desirable characters is the key towards developing successful hybrids. Existence of genetic diversity among the inbred lines provides an opportunity for plant breeders to develop hybrids with desirable features. The main issue of hybrid breeding is that it utilizes heterosis which is mainly related to the genetic distance among inbred lines, so, further relatives exhibit higher heterosis. Therefore, a plant breeder should know the relatedness or genetic relationship of inbred lines to obtain higher yield which is the goal (Hladni *et al.*, 2008).

Principle component analysis has been proved to be a very useful statistical tool in identifying genotypes with superior performance and classifying lines based on genetic distance in many crops like sunflower (Rehman *et al.*, 2012; Iqbal *et al.*, 2013; Hussain *et al.*, 2017; Venujayakanth *et al.*, 2017), wheat (Hailu *et al.*, 2006), black gram (Ghafoor and Arshad, 2005). Principle component analysis also helps to determine inbred lines which can be used as parents to improve sunflower seed yield and oil contents (Kholgi *et al.*, 2011). This study was organized with the objective of assessing variability among sunflower inbred lines at genetic level and to utilize the outcomes in future sunflower improvement programs.

MATERIALS AND METHODS

During spring, 2018, 24 Sunflower inbred lines were planted thrice in a randomized complete block design at the research farm of Oilseeds Research Institute, Faisalabad (Pakistan). The inbred lines with prefix "ORI" represent the cytoplasmic male sterile lines (A lines) whereas prefix "RL" represents the fertility restorer (R lines). The data were noted down from 10 arbitrarily selected plants from each replication. Standard agronomic and plant protection practices were carried out during the conduct of whole experiment. Recommended dose of NPK (118:85:62 kg/ha) was applied. The field was irrigated 5 times due to the extreme weather conditions during critical growth stages. After the completion of physiological maturity (100–110 days) data were noted for head diameter (HD), days to maturity (DTM), achene yield per plant (AYP), oil contents OC), oleic acid (OA) and linoleic acid (LA). Oil percentage was determined through Near Infra-Red Spectroscopy (NIR) at Oil technology laboratory, Oilseeds Research Institute, Faisalabad (Pakistan).

Table No.1: List of twenty-four inbred lines of sunflower developed by Oilseeds Research Institute, Faisalabad (Pakistan).

Sr. No.	Inbred lines	Sr. No.	Inbred lines
1	ORI-2	13	ORI-95
2	ORI-7	14	ORI-99
3	ORI-10	15	ORI-100
4	ORI-19	16	RL-38
5	ORI-20	17	RL-39
6	ORI-21	18	RL-50
7	ORI-22	19	RL-66
8	ORI-23	20	RL-67
9	ORI-25	21	RL-68
10	ORI-26	22	RL-86
11	ORI-92	23	RL-100
12	ORI-94	24	RL-109

STATISTICAL ANALYSIS:

Data noted for various characters were investigated through analysis of variance (Steel *et al.*, 1997) to assess the presence of useful genetic variability by using statistics 8.1 and genotypes were selected through GGE biplot analysis (Yan *et al.*, 2000) by using XLSTAT version of Microsoft Excel, Copyright Addinsoft 1995-2012 (<http://www.xlstat.com>). LSD test at P = 0.05 is used to access the association among treatments for yield or yield components. Principle component analysis is used to calculate the mathematical analysis or simple statistical analysis.

RESULTS

Variance analysis displayed that sunflower inbred lines had evidently variable performance for characters associated with seed yield and oil quality (Table No. 2). Means of studied traits (Table No. 3) showed that inbred line ORI-92 takes maximum number of days to reach its maturity stage and had highest HD. Among all inbred lines, ORI-25 and ORI-94 displayed uppermost AYP. Inbred line RL-67 had highest percentage of OC followed by ORI-19. Maximum percentage of OA was exhibited by RL-50 whereas lowest OA was presented by ORI-100. Among R lines, RL-86 had highest LA percentage whereas lowest was displayed by RL-100. The principal components having more than 1 eigen value are presented in Table No. 4. First two PCs contribute 62% in total variability. First principle component points out the importance of HD, AYP and LA and suggested the positive association between these variables. Inbred lines that fall in PC1 vary together for observed characters. Second principle component is associated with DTM and OC. These

variables have negative association between them. Similar results have been displayed in graphical presentation of bi-plot analysis (Fig. 1) in which PC1 is laid against PC2. Length and direction of vectors in Fig. 1 represents the extent and direction of variability among characters under investigation. Plotting of various inbred lines in four quadrates displayed their worth in terms of genetic diversity present among them and helps in their selection for inclusion in hybridization program for getting desirable recombinants. Inbred lines that are plotted along the vectors associated with various characters had high degree of variability for that character. Inbred lines

ORI-94 and ORI-25 showed its worth for AYP, HD and OC whereas the inbred line ORI-92 have good percentage of LA and had long maturity duration. RL-68, RL-67 and RL-66 displayed their value for OA content. Associations among various characters are very much important to design selection criterion for the development of desirable cross combinations. Substantially positive association was observed between AYP, HD and LA whereas negative relationship was observed for these characters with OA. LA had negative association with OA (Qamar *et al.*, 2018).

Table No. 2: Mean squares for yield and oil quality related parameters in sunflower inbred lines.

SOV	DF	HD	DTM	AYP	OC	OA	LA
Replications	2	7.063	0.0007	98.13	0.148	8.0	14.79
Inbred lines	23	47.75	55.54	267.5	38.20	420.3	524.2
Error	46	3.81	0.0001	49.58	4.83	9.96	14.66
Total	71						

SOV = Source of variation, DF = degrees of freedom, HD = Head diameter, DTM = days to maturity, AYP = Achene yield per plant, OC = Oil contents, OA = Oleic acid, LA = Linoleic acid

Table No. 3: Mean values for achene yield and oil related characters in sunflower inbred lines

Sr. No.	Inbred line	HD	DTM	AYP	OC	OA	LA
1	ORI-2	14.58	99.00	19.20	34.70	61.31	39.31
2	ORI-7	17.25	99.00	24.65	35.09	40.00	42.14
3	ORI-10	9.50	96.00	20.37	38.59	38.61	56.46
4	ORI-19	12.50	96.00	25.55	43.12	40.09	36.94
5	ORI-20	14.33	96.00	26.28	40.45	45.96	44.34
6	ORI-21	17.33	98.00	30.42	40.57	44.33	42.93
7	ORI-22	12.50	96.00	23.52	42.64	49.53	41.55
8	ORI-23	13.33	95.00	11.65	33.02	43.38	41.46
9	ORI-25	16.83	94.00	35.64	40.17	40.90	40.62
10	ORI-26	16.00	98.00	25.07	42.59	47.78	43.46
11	ORI-92	17.50	112.00	25.28	35.02	37.07	55.86
12	ORI-94	13.67	97.00	35.46	39.21	35.84	59.30
13	ORI-95	13.33	103.00	7.21	35.76	35.10	29.49
14	ORI-99	14.33	103.00	26.44	39.71	43.27	34.37
15	ORI-100	17.00	95.00	25.24	39.60	34.31	43.79
16	RL-38	13.00	101.00	14.62	37.92	66.28	29.82
17	RL-39	14.00	97.00	28.10	36.60	66.55	23.87
18	RL-50	12.00	98.00	22.82	40.51	71.61	16.87
19	RL-66	7.00	98.00	8.51	37.17	63.84	25.25
20	RL-67	5.67	100.00	8.00	45.37	51.57	15.12
21	RL-68	6.00	91.00	4.57	36.13	63.60	24.77
22	RL-86	5.67	98.00	7.35	33.33	61.01	59.70
23	RL-100	5.33	96.00	5.83	31.06	36.59	13.69
24	RL-109	13.67	106.00	15.41	35.83	56.73	40.92

HD = Head diameter, DTM = days to maturity, AYP = Achene yield per plant, OC = Oil contents, OA = Oleic acid, LA = Linoleic acid

CONCLUSION

Analysis of variance suggested the presence of variation in available germplasm. ORI-94, ORI-92 and ORI-25 displayed its value for AYP, HD and LA whereas RL-50, RL-68 and RL-66 is worthy for it OA content. Cytoplasmic male sterile lines and fertility restorer lines that falls in opposite quadrates are important in terms of hybrid development program as genetically divergent parents are prerequisite for development of high-profile hybrids. Moreover, relationship among characters showed that AYP can be improved by improving HD. Among quality factors, both important fatty acids like LA and OA cannot be improved simultaneously.

References

- [1]. Aftab, M., T. Mahmood and H.S.B. Mustafa. 2019. Prospects of Oilseeds in Pakistan, Directorate of Oilseeds, Ayub Agricultural Research Institute, Faisalabad (Pakistan) ISBN:978-969-7594-14-6.
- [2]. Aguirrezabal, L., P. Martre, G. Pereyra-Irujo, M.M. Echarte and N. Izquierdo. 2015. Improving grain quality. In: Eco physiological and Modeling Tools to Develop Management and Breeding Strategies, pp: 423–465. Victor, S. and C. Daniel (eds.). Crop Physiology. Lamber Academic Publ., Saarbrucken, Germany.
- [3]. Ali, M.A., N.N. Nawab, A. Abbas, M. Zulkiffal and M. Sajjad. 2009. Evaluation of selection criteria in *Cicer arietinum* L. Using correlation coefficients and path analysis. Aust. J. Crop Sci., 3: 65–70.
- [4]. Arshad, M., M.K. Ilyas and M.A. Khan. 2007. Genetic divergence and path coefficient analysis for seed yield traits in sunflower (*Helianthus annuus* L.) hybrids. Pak. J. Bot. 39(6): 2009-2015.
- [5]. Economic Survey of Pakistan, 2018-19. Finance Division Government of Pakistan.
- [6]. Ghafoor, A. and M. Arshad. 2008. Multivariate analysis for quantitative traits to determine genetic diversity of blackgram [*Vigna mungo* (L.) Hepper] germplasm. Pak. J. Bot. 40(6): 2307-2313.
- [7]. Hailu, F., A. Merker, H. Singh, G. Belay and E. Johansson. 2006. Multivariate analysis of diversity of tetraploid wheat germplasm from Ethiopia. Genet. Resour. Crop Evol. 53: 1089-1098.
- [8]. Hladni, N., S. Jocić, V. Miklič, A. Mijić and D.S. Panković. 2014. Direct and indirect effects of morphological traits on seed yield of sunflower (*Helianthus annuus* L.). DOI: <https://doi.org/10.2298/hel1053101h>
- [9]. Hussain, F., M. Rafiq, M. Ghias, R. Qamar, M.K. Razzaq, A. Hameed, S. Habib and H.S.B. Mustafa. 2017. Genetic Diversity for Seed Yield and its Components Using Principal Component and Cluster Analysis in Sunflower (*Helianthus annuus* L.). Life Sci. J. 14(5): 71-78.
- [10]. Iqbal, M., U. Ijaz, Samiullah, M. Iqbal, K. Mahmood, M. Najeebullah, Abdullah, S. Niaz and H.A. Sadaqat. 2013. Genetic divergence and path coefficient analysis for yield related attributes in sunflower (*Helianthus annuus* L.) under less water conditions at productive phase. Plant Knowled. J. 2(1): 20-23.
- [11]. Khan, H., S. Muhammad, R. Shah and N. Iqbal. 2007. Genetic analysis of yield and some yield components in sunflower. Sarhad J. Agric. 23(4): 987-990.
- [12]. Kholgi, M., I. Bernousi, R. Darvishzadeh and A. Pirzid. 2011. Correlation and path-coefficient analysis of seed yield and yield related trait in Iranian confectionary sunflower population. Afr. J. Biotechnol. 10(61): 13058-13063.
- [13]. Kholgi, M., I. Bernousi, R. Darvishzadeh, A. Pirzad and H.H. Maleki. 2011. Collection, evaluation and classification of Iranian confectionary sunflower (*Helianthus annuus* L.) populations using multivariate statistical techniques. Afr. J. Biotechnol. 10(28): 5444-5451.
- [14]. Mustafa, H.S.B., J. Farooq, E. Hassan, T. Bibi and T. Mahmood. 2015. Cluster and Principle Component Analysis of Maize Accessions under Normal and Water Stress Conditions. J. Agric. Sci. 60(1): 33-48.
- [15]. Neto, A. R., A. M. R. O. Miguel, A. L. Mourad, E. A. Henriques and R. M. V. Alves, 2016. Environmental effect on sunflower oil quality. Crop Breed. Appl. Biotechnol., 16: 197–204
- [16]. Qamar, R., A. Bibi, H.A. Sadaqat, F.S. Awan and H.M. Akram. 2018. Computation of Differential Response of Sunflower Genotypes for Achene Yield and Oil Quality against Lead Toxicity. Int. J. Agric. Biol. 20(12): 2731-2736.
- [17]. Ram, J.J., U.K. Singh, S.K. Singh and B. Krishna. 2018. Study of genetic diversity in

- sunflower (*Helianthus annuus* L.). Int. J. Curr. Microbiol. App. Sci. 7(5): 2266-2272.
- [18]. Rehman, R., M. Arshad, M.A. Khan, A.S. Mohmad, G. Shabbir and M.K.N. Shah. 2012. Using Multivariate Analysis for Selecting Desirable Hybrids in Sunflower (*Helianthus annuus* L.). Pak. J. Bot. 44(5): 1715-1720.
- [19]. Steel, R.G.D. 1997. Principle and Procedures of Statistics. McGraw Hill Book Company Inc. New York, USA.
- [20]. Yan, W., L.A. Hunt, Q. Sheng, and Z. Szlavnic. 2000. Cultivar Evaluation and Mega-Environment Investigation Based on the GGE Biplot. Crop Sci. 40:597-605. doi:10.2135/cropsci2000.403597x.

6/2/2022