



## The effect of Acetyl-cysteine and carnitine on post-thaw sperm characteristics of the Sand Gazelle (*Gazella subgutturosa marica*)

Mohammed Al-zharani

Biology Department, College of Science, Imam Mohammad ibn Saud Islamic University (IMSIU), Riyadh 11623, Kingdom of Saudi Arabia

### Abstract

The current study aimed to estimate the effect of antioxidants (N-acetyl-cysteine (NAC) and L-carnitine (LC)) as an antioxidant added on triladyl semen extender of gazelles. The results showed no significant effect on progressive and mass motility characteristics when compared to the control at ( $P > 0.05$ ). Assessment of acrosome abnormality and membrane integrity were assessed for (NAC) and (LC), the result was  $2.40 \pm 0.12\%$  for (NAC) for acrosome abnormality and  $49.40 \pm 3.06\%$  for membrane integrity, whereas the (LC) was ( $2.40 \pm 0.16\%$  for acrosome abnormality and  $42.60 \pm 2.94\%$  for membrane integrity respectively with no significant differences among groups. on the other hand, the viability was ( $3.20 \pm 0.17\%$ ) for triladyl with (NAC), and ( $3.42 \pm 0.42\%$ ) for (LC) added to triladyl, furthermore, the (NAC) reduced DNA damage when compared to control group ( $3.20 \pm 0.17\%$ ) at ( $P > 0.05$ ). Among all groups, the pregnancy rates do not show significant differences.

[Mohammed Al-zharani . **The effect of Acetyl-cysteine and carnitine on post-thaw sperm characteristics of the Sand Gazelle (*Gazella subgutturosa marica*)**. *Life Sci J* 2022;19(12):19-26]. ISSN 1097-8135 (print); ISSN 2372-613X (online). <http://www.lifesciencesite.com>. 03.doi:[10.7537/marslsj191222.03](https://doi.org/10.7537/marslsj191222.03).

**Keywords:** antioxidants, Gazelle, cysteine, carnitine, motility, acrosome, membrane integrity, DNA damage.

### Introduction

Sand gazelle (*Gazella subgutturosa marica*) Population and distribution has significantly affected due to the degradation of natural habitats and excessive hunting during the last decade, so that they are living in a small range and in a few isolated groups as a protected area. Saudi Wildlife Authority has research - breeding centers for specific species, one in all them is that the sand gazelle. A technique of importance in the genetic management of endangered species is the cryopreservation of sperm, it's a regular procedure used in a variety of state inclusive assisted reproduction technologies for human and animals and recently for small ruminants and endangered species such as gazelles (Garde J, *et al* , 2007), nevertheless, sperms are undergoing to effective changes in temperature, the formation of the ice crystal, and various stresses such as (physical, chemical, diffusion, and oxidative) throughout the cryopreservation method, that solely compromise spermatozoon fertility and quality (Ezzati M *et al* 2020 and Watson P, *et al* , 1995), Previous studies by (Woolley D and Richardson D, 1978) showed that its cause changes in sperm morphology and effect on motility as reported by (Henry M *et al* , 1993), furthermore the membrane extremely sensitive for freezing and cause cryodamage as cold shock and formation of ice crystal (Aitken R, *et al* , 1989) and (Alvarez J and Storey B, 1995), the membranes can

be expose to various stresses as osmotic variations, lipid and protein restructurings, oxidative stress (Bailey J, *et al* , 2000) and (Watson P, 1995). Oxidative stress has been known in concert various intermediary male physiological state by inflicting sperm dysfunction. at normal condition the reactive oxygen species (ROS) are composed due to reduction of oxygen, therefore as a result sperms loss their motility, integrity and fertility (Aitken R J, Fisher H, 1994) and (Gandini L *et al*, 2000). Due to the richness of cell membranes with polyunsaturated fatty acids this increase the sensibility of sperms to oxidative stress, that produce (ROS) (Bansal A, *et al*, 2010 and Gharagozloo P, *et al* 2011). highly concentration of polyunsaturated requires effectual antioxidant to preserve against peroxidation and dysfunction associated with sperm (Aitken R, *et al*, 1989) and (Alvarez J and Storey B, 1995). sperms and seminal plasma have an antioxidant to avoid oxidative stresses. Nevertheless, the capacity of the cells very limited due to the limitation on antioxidants contents to scavenge oxidants (Storey B, 1997). Antioxidant component properties reduce the effect of damage from ROS and cold shocks (Fardin A, *et al*, 2016), (Alvarez J and Storey B, 1983) and (Donnelly E, *et al* 2000), therefore to improve sperm quality for human (Alvarez J and Storey B, 1989), ram (Aisen E, *et al* 2002), canine (Michael A , *et al* , 2007), (Bucak M,

*et al* , 2008), and goat (Bucak M, *et al* , 2009).the derivatives of amino acid L-cysteine containing sulfur-containing amino acids named thiol compounds are currently used mainly as an antioxidant and cytoprotective in mammalian cells. (Zafarullah M, 2003), In addition, daily treatment with NAC results in a significant improvement in sperm motility (Safarinejad M and Safarinejad S, 2008). Even though (NAC) lowering endogenous oxidant levels and to maintain cells against harmful stress, however, it's a poor scavenger and the molecular mechanisms have remained uncertain (Daria E *et al*, 2018). Furthermore, (Ciftci H, *et al*, 2009) and (Oeda T, *et al* 1997) reported that (NAC) has free radical scavenging activity both in vivo and in vitro respectively. the post-thaw characteristics for bull (Sariozkan S, *et al*, 2009), ram (Uysal O and Bucak M, 2007), and goat sperm (Bucak M and Uysal O, 2008) were improved by added of (NAC), moreover, its, maintain the viability, chromatin structure and membrane integrity of boar sperm during liquid storage at 5 °C for goat sperm (Szczesniak B, *et al*, 2003). On the other hand, L-carnitine (LC) highly concentrated in the epididymis of sperms, and has a vital role in the production of energy by transferring fatty acids into mitochondria, its transport long chain fatty acids from cytosol into the mitochondria matrix and play critical role in the regulation bioenergetic metabolism and respiration in most spices including gazelle as published by (Aleissa M, *et al* , 2007) and (Mongioi L, *et al*, 2016) , moreover, it is protects the cell, mitochondrial membrane, and DNA integrity against free radicals (Adewoyin M, *et al*, 2017), also its enhancement of sperm fertility, count, and motility (Tahereh M, *et al* , 2019). many studies prove that a close interrelationship among semen carnitine (LC) together with attributes of semen moreover, these studies were confirmed amongst diverse species such as of ram, rats, stallions and human (Brooks D, 1979), stallions (Stradaioli G, *et al*, 2004).

Semen diluent composition is a highly important factor influencing freezing ability. The role of extenders is to protect sperms from cold shock and agglutination; maintain stable pH and osmotic pressure. as far as we know, the antioxidant contributes to protect cells against cryodamage, but due to lack an information and study regarding sand gazelle sperm cryopreservation and influences of (NAC) and (LC). currant study designed to carry out an influence adding N-acetyl-cysteine (NAC) and (LC) into triladyl extender and study the effect of them on characteristics of frozen thawing sperms, viability and fertilizable.

## Methodology

### Gazelles preparation:

The commercially available diluent (Triladyl by Minitube, Tiefenbach, Germany), was used in this study for the Cryopreservation of semen. The antioxidants, carnitine, lysine and antibodies were obtained from Sigma–Aldrich Chemical Co, USA. Study was conducted in The King Khalid Wildlife Research Centre (KKWRC), Saudi Arabia. 10 male of the sand gazelles (*Gazella subgutturosa marica*), average weight was 15-18 kg and captured manually by using canvas trap around food point, gazelle's heads were immediately covered after captured to reduce the stress. By using the electroejaculation method semen was collected according to the procedure described by (Al-Eissa M, *et al*, 2007) based on the method published by (Holt W, *et al*.1996). Prior collection the gazelles were sedative by using xylazine (8.4 ± 1.5 mg/kg) and ketamine hydrochloride (6.9 ± 1.5 mg/kg). 120 ejaculates were collected from the gazelle (12 ejaculates for each gazelle), semen was collected in a conical tube then immediately transfer to water bath at 37C and evaluated within 30 to 60 min of collection. Samples with a sperm concentration of more than  $2.8 \times 10^9$  /ml, and least 80% motility were used for CASA. Ejaculated is measured by graduated pipette and sperm count by photometer sperma cue (model 12300/0500, Minitube, Germany). 15% of egg yolk and 6% glycerol (v/v) were added to triladyl containing N-acetyl-cysteine (NAC), (LC), then pH was adjusted to 6.8. Each ejaculate was divided into 3 portions then dilute to  $60 \times 10^6$ /ml sperms. 0.25ml French straws were used to load diluted semen samples then frozen by digital freezing machine (Freeze control, CI-8000, from Crtologic pty.ltd) afterward, then transferred to vapor nitrogen for 24 hours later on, samples were thawed in water bath for 20s. at 38 °C

### Sperm motilities Assessment

Sperm velocity parameters by computer-assisted sperm analyzer (CASA) such as Hamilton-Thorne IVOS 11, Biosciences, Beverly, MA 01915 USA). IVOS CASA system yields precise, objective assessment, and relabels result in different semen parameters based on the estimating of individual sperm (Perumal P, *et al*, 2014). Recent reports by (Mortimer S, *et al*, 2015), proposed that, CASA does not estimate the proportionality of sperms motile only but also estimates other sperm velocity. the system is equipped with a high-resolution camera connected to a phase-contrast microscope moreover it is needs standardization and validation concerning semen preparation, calibration,

and technical settings to provide precise results. For each evaluation 5 µl of diluted sample was placed in the sperm chamber and up to 15 microscopic fields were analyzed to include at least 100 sperms. CASA system is able to measure many sperm velocity include curvilinear velocity (VCL), straight linear velocity (VSL), mean velocity (VAP), linear coefficient (LIN) and amplitude of lateral head displacement (ALH)

### Sperm abnormalities Assessment

According to methods published by (Sekoni *et al.* 1988), morphology of sperms was examined by using hancock Stain (P/N HCS-101), the staining mixture consisted of 5% of nigrosin and 1% eosin B in 3% sodium citrate dehydrates solution. drop of semen was added to drop of the stain, then mixed gently, a fresh smear was applied on slide. At least, 100 sperms, live -dead sperms in an ejaculate were counted and estimated by a microscope at magnification 40x.

### Integrity membrane evaluation

Evaluation of membrane integrity carried out by using HOS test according to the method described by (Jeyendran R, 1984), 10 ml of semen with 20 ml of hypo osmolar solution were examined by counting 200 sperm at magnification 40x. Sperm with swollen tail was consider as positively to HOS test (Buckett W, *et al.*, 1997).

### DNA viability assessment

According to the method was described by (Yaniz J *et al.*, 2013) the DNA viability was estimated by a commercial kit contain acridine orange- propidium iodide, 10 ml of semen with 20 ml of stain applied on glass slide then by using fluorescence microscope the viability was estimated according to the procedure results in selective labelling of live and dead cells visualized in green and red fluorescence colors respectively. A total of 200 sperms were assessed/sample.

### Fertility rate

The evaluation of sperm, morphology and quality are highly significant during an evaluation of male fertility due to the strong correlation between them. an accurate morphological assessment for each individual sperm (Varner, D, 2016), 10 female gazelles were artificially inseminated (AI) with frozen-thawed semen. The ability of sperm for fertilizing was estimated based on the non-return rates (proportion %) by the formula given below, the

effective non-return rates (NNR) was determined at 80 days post-insemination.

$$NNR = \frac{\text{total no of females inseminated} - \text{no of females returning to estrus}}{\text{total no of females inseminated}} * 100$$

Table 1. Mean (±SE) of post-freeze-thawing sperm motility for Gazelle semen supplemented with different antioxidants.

Groups	Progressive motility (%)	Mass motility (%)
Control	59.50±2.14	68.32±3.55
(LC)	57.21. ±2.4	69.22±2.14
(NAC)	57.44 ±1.3	71.54±6.22

All values are expressed as mean ± SE, Different superscripts in the same column demonstrate significant differences(P> 0.05).

### Statistical analysis

Statistical analyses were performed by using the SPSS 17 package program. Results were expressed and as mean ± SEM for all sperm motility, velocity characteristics, abnormalities, and (HOST) tests by analysis of variance significant differences between the groups. The analysis of variance used was compared for DNA damage between groups. Pearson's chi-square test was used to compare the pregnancy rate.

### Results

The antioxidant added to diluent in this experiment do not reflect any significant on sperms progressive or, motility characteristics by (CASA) when compare to control group at (P > 0.05) as show in table 1, (57.21. ±2.4%) for (LC) and (57.44 ±1.3) for (NAC) for Progressive motility and furthermore (69.22±2.14), (71.54±6.22) for mass motility. Data of sperm characteristic's by CASA as showed in table 2 are (VAP, VSL and VCL, ALH and LIN) were. the motility pattern parameter for VAP was (114.4±2.1), (118.34±42) for (LC), (NAC) respectively and (119.4±1.2) for control group with no significant difference among them. There were no significant differences in the (VSL) for all treatment compare to control they were (76.26±4.19) for (LC), (77.46±3.6) for (NAC) and (80.12±2.0) for control. None of the (VCL) values were significantly different between groups even (LC) (125.3±6.7), (NAC) (143.6±1.7) or control (134.6±2.4). furthermore, the (ALH) also shows that, the (LC) and (NAC) values were

(10.3±2.8) and (12.4±1.4) respectively and significantly different to the control (22.4±0.24) at (P> 0.05). Finally, linearity (LIN) is the linearity of the curvilinear trajectory calculated as (VSL/VCL) × 100, according to table 2 we don't show and significant between three groups (LC) was (18.4±4.3), (NAC) was (16.62±4.3) and control was (17.34±6.2).

Table 2. Mean (±SE) CASA determination of sperm velocity in post-freeze-thawing sperm of Gazelle semen supplemented with different antioxidants.

Group	VAP	VSL	VCL	ALH	LIN
Control	119.4 ±1.2	80.12±2.0	134.6 ±2.4	22.4±0.24	17.34 ±6.2
(LC)	114.4 ±2.1	76.26±4.19	125.3 ±6.7	10.3±2.8 <sup>a</sup>	18.4±4.3
(NAC)	118.3 ±4.2	77.46±3.6	143.6 ±1.7	12.4±1.4 <sup>b</sup>	16.62 ±4.3

All values are expressed as mean ± SE, values with different superscripts in the same column demonstrate significant differences at (P> 0.05). VAP=Average value (mm/s), VSL=Rectilinear speed (mm/s), VCL=Curvilinear velocity (mm/s), ALH=, Amplitude of lateral head (mm) and LIN=Linearity index (%).

Semen extender provided with (NAC) and (LC) give a result for acrosomal abnormality as (2.60% ± 0.24) and (2.40% ± 0.16) individually, lower than control and with a significant difference at (P > 0.05) compare to control 8.00% ± 0.44. Regarding the integrity of membrane There were no significant differences between (NAC), (LC) (49.40 ± 3.06%) and (42.60 ± 2.94%), respectively) compare to control (41.40±4.74) but it's clearly that the treatment with (NAC) had more positive effect on the membrane integrity when compare with others. Moreover, the statistical analysis for data collected for viability showed, there are no significant difference between the two groups although the treatment groups were reduced viability compare with control at (P > 0.05) as seen in table 3. Pregnancy rates show no significant difference at (P > 0.05) between the diluent groups (NAC) was 44%, (LC) was 50% and control 47% as showed in table 4.

Table 3. Mean (±SE) for Acrosomal abnormality, integrity of membrane and viability of post-freeze-thawing sperm of Gazelle semen supplemented with different antioxidants.

Group	Acrosomal abnormality (%)	integrity of membrane (%)	viability (%)
Control	8.00 ± 0.44	41.40±4.74	4.19 ± 0.24
(LC)	2.40 ± 0.16*	42.60±2.94	3.42±0.42
(NAC)	2.60 ± 0.12*	49.40±3.06	3.20 ± 0.17

All values are expressed as mean ± SE, Different superscripts in the same column demonstrate significant differences(P> 0.05).

All values are expressed as mean ± SE, Different

Table 4. Pregnancy rates after artificial insemination with freeze-thawing semen of gazelle.

Groups	Pregnancy rate (%)
Control	47%
(LC)	50%
(NAC)	44%

superscripts in the same column demonstrate significant differences(P> 0.05).

### Discussion and conclusion

Sperms Motility reflects the ability of a sperm to transfer through the female genital tracks and acquire ability to penetrate several ovum membranes and fertilizing it. A positive correlation between sperm motility and fertilizing capacity has been shown in many species (Steven P. *et al* 2011). In this study we examined the influence of (NAC) and (LC) as an antioxidant on sperm characteristics and oxidative stress characteristics post freeze–thawing of gazelle's semen, our results were agreement with the study achieved by (Bucak M, *et al* ,2010a). From the previous result, it can be realized that, motility characteristics do not have sufficient information for the sperms quality and don't showed differences between the effect of (NAC) and (LC) as highlighted in Table 2,3. Comparison of the differences between (NAC) and (LC) the mean values of the Acrosomal abnormality was lower than control and with a significant difference at (P > 0.05) compare to control, membrane integrity within (NAC) and (LC) revealed that there were no significant changes between them compare to control. our results show similarities with

the previous studies, reported that the sperm parameters do not show significant relationship between antioxidant (LC) therapy and semen parameters (Sedigheh A, *et al*, 2016). This study show that, the triladyl with antioxidants was very efficient in preservation of mass motility and integrity of plasma membrane ( Vera.O *et al* , 2009), also the antioxidant, such as (NAC) and (LC) can effectively improve semen parameters, it was similar to studies done by (Sedigheh A, *et al* 2016) for men and bovine (Bucak M *et al* ,2010a), goat (Bucak M, *et al* , 2010b) against cryodamage. (NAC) and (LC) may equalize free radicals and protects sperm membrane against stress (Mazor D *et al* 1996). In addition, structure of the plasma membrane is consisting of high levels of polyunsaturated fatty acids (PUFAs) that improve of membrane flexibility and makes it susceptible to attacked by ROS (Eskenazi B, *et al*, 2005) and (Sheweita S, *et al*, 2005). and eventually loss in fertility (Alvarez J *et al*, 1993) and (Griveau J *et al*, 1995). Cryoprotectants are improved recoveries of cells after cryopreservation and can assist in reducing stress effects. Recently many studies focused mainly on the different additives in the diluent in order to acquire the higher post-thaw motility and fertility rates. Supplementation with (NAC) and (LC) to the cryoprotectants contributes in the protection of DNA materials against cryodamage, In the present study, our result was similar with previous study mentioned by (Bucak M, *et al*.,2010a). Recent studies showed that long-term exposure of sperm to long-term sperm storage caused a theatrical damage in sperm characteristics, this study aims to focus on importance of antioxidative and its role as cryoprotectant. During the (HOS) test, sperm with intact plasma membranes go through swelling due to the inflow of water and thus, increase in volume to institute equilibrium between the extra- and intracellular compartments, the test was applied to goat (Ranjan R, *et al*. 2009). Freezing and long-term storage makes sperm plasma membrane capable to loss its integrity then function, resulting from reactive oxygen species. Viability HOST response showed clearly that, (NAC) and (LC) don't have any protective effect to protect membrane during cryopreservation and remains to be examined its parallel to result for ram by (Aisen E, *et al* 2002). Incubation of the sperms by using fluorochromes is considered important prior to the assessment of membrane integrity as reported by (Pintado B, *et al*, 2000; Brito L, *et al*., 2003; Yaniz J, *et al*., 2008 and Yaniz J, *et al*, 2012). since the AO penetrating live sperm and bound to DNA while the PI penetrating the damaged sperm and replacement of

AO. our result showed that, the antioxidants (NAC), (LC) unable to maintain DNA integrity, these results was different than study reported on bovine sperm by (Bucak M, *et al* , 2010A), for the signify role of (NAC) and (LC) in the maintenance of DNA integrity. the diluent contains scavengers that are appropriate to maintain sperms against oxidative stress during cryopreservation, moreover, new findings indicated for no correlation between improvement fertility and supplemented to the diluent, and this finding with our results (Sariozkan S, 2014). The following conclusions can be drawn from the present study antioxidants enhanced sperm abnormalities and acrosomal damage for thawed sample. Also, (NAC), (LC) providing an improved DNA integrity. Further research should be done to investigate the effects of different antioxidant on cryopreservation and fertility of endanger Species specially gazelle of the Arabian Peninsula due to Scarcity of information about them.

## References

- [1]. Adewoyin M., Ibrahim M., Roszaman R., Isa M., Alewi N., Rafa A., Anuar M. (2017). Male Infertility: The Effect of Natural Antioxidants and Phytocompounds on Seminal Oxidative Stress. *Diseases*. 5(1):9. 2-26.
- [2]. Aisen E.G, Medina V.H, Venturino.A. (2002). Cryopreservation and post-thawed fertility of ram semen frozen in different trehalose cocentrations. *Therionology*. 15;57(7):1801-8.
- [3]. Aitken R J, Fisher H. (1994). Reactive oxygen species generation and human spermatozoa: the balance of benefit and risk, *BioEssays* 16, 259–267.
- [4]. Aitken R J, Clarkson J S, Fishel S. (1989). Generation of reactive oxygen species, lipid peroxidation, and humansperm function, *Biol. Reprod*. 40 183–197.
- [5]. Aleissa M and Alhomida A. (2007). A study of the distribution of total, free, short-chain acyl and long-Chain acyl carnitine in whole blood and plasma of arabian sand gazelles (*Gazella subgutturosa marica*). *Comparative Haematology International*. 1:65-69
- [6]. Al-Eissa M, AlHamidi A and Kandeal S,( 2007). Semen cryopreservation using Triladyl and Tris diluents of the Arabian and Gazelle Males (*Gazella subgutturosa marica*). *Arab Gulf Journal of Scientific Research* 25(4):199-206.
- [7]. Alvarez J G and Storey BT. (1983). Taurine, hypotaurine, epinephrine and albumin inhibit lipid peroxidation in rabbit spermatozoa and

- protect against loss of motility, *Biol. Reprod.* 29, 548–555.
- [8]. Alvarez J G and Storey B T. (1989). Role of glutathione peroxidase in protecting mammalian antioxidant enzymes: their source, function in the female, and ability to preserve sperm DNA integrity in the golden hamster, *J. Androl.* 24, 704–711.
- [9]. Alvarez J, Storey B. (1995). Differential incorporation of fatty acids into and peroxidative loss of fatty acids from phospholipids of human spermatozoa, *Mol. Reprod. Dev.* 42 334–346.
- [10]. Bailey J L, Bilodeau J F, Cormier N. (2000). Semen cryopreservation in domestic animals: a damaging and capacitating phenomenon, *J. Androl.* 21, 1–7.
- [11]. Bansal AK, Bilaspuri G. (2010). Impacts of oxidative stress and antioxidants on semen functions. *Vet Med Int*, 686137.
- [12]. Brooks D E. (1979). Carnitine, acetylcarnitine and the activity of carnitine acyltransferases in seminal plasma and spermatozoa of men, rams and rats, *J. Reprod. Fert.* 56 667–773.
- [13]. Brito L, Barth A, Bilodeau, Goeseels S, Panich PL, Kastelic J. (2003). Comparison of methods to evaluate the plasmalemma of bovine sperm and their relationship with in vitro fertilization rate. *Theriogenology* 60: 1539-1551
- [14]. Bucak M.N, Atessahin A, Yuce A. (2008). Effect of anti-oxidants and oxidative stress parameters on ram semen after the freeze-thawing process, *Small Rumin. Res.* 75, 128–134.
- [15]. Bucak M N and Uysal O. (2008). The role of antioxidants in the freezing of Saanen goat semen, *J. Indian Vet.* 85 (2008) 148–150.
- [16]. Bucak M N, Sariozkan S, Tuncer P B, Ulutas P A, Akcadag HA. (2009). Effects of antioxidants on microscopic semen parameters, lipid peroxidation and antioxidant activities in Angora goat semen following cryopreservation, *Rumin. Res.* 81, 90–95.
- [17]. Bucak M, Sariozkan S, Tuncer P, Sakin F, Atessahin A, Kulaksiz R, Cevik M. (2010b). The effect of antioxidants on post-thawed Angora goat (*Capra hircus ancyrensis*) sperm parameters, lipid peroxidation and antioxidant activities, *Small Ruminant Res.* 89, 24–30.
- [18]. Buckett W, Luckas M, Aird I, Farquharson R, Kingsland C, Lewis-Jones D. (1997). The hypo-osmotic swelling test in recurrent miscarriage, *Fertil. Steril.* 68, 506–509.
- [19]. Bucak M , Tuncer P , Serpil S , Başpınar N, Taşpınar M, Cayan K, Bilgili A, Peker P, Akalin P, Buyukleblebici S, Aydos S, Ilgaz S, Sunguroglu A, Oztuna D. (2010A). Effects of antioxidants on post-thawed bovine sperm and oxidative stress parameters: antioxidants protect DNA integrity against cryodamage, *Cryobiology* 61(3), 248–253.
- [20]. Ciftci H, Verit A, Savas M, Yeni E, Erel O. (2009). Effects of N-acetylcysteine on semen parameters and oxidative/antioxidant status. *Urology.* 74(1):73–76.
- [21]. Daria Ezeriņa, Yoko Takano, Kenjiro Hanaoka, Yasuteru Urano, and Tobias P. Dickl. (2018). N-acetyl cysteine functions as a fast-acting antioxidant by triggering intracellular H<sub>2</sub>S and sulfane sulfur production. *Journal Cell chemical biology.* 25, (4), 19, Pages 447-459.
- [22]. Donnelly E T, McClure N, Lewis SE. (2000). Glutathione and hypotaurine in vitro: effects on human sperm motility, DNA integrity and production of reactive oxygen species, *Mutagenesis* 15, 61–68.
- [23]. Eskenazi B, Kidd S, Marks A, Slotter E, Block G, Wyrobek A. (2005). Antioxidant intake is associated with semen quality in healthy men. *Hum Reprod.* 20: 1006-1012.
- [24]. Ezzati, M.; Shانهbandi, D.; Hamdi, K.; Rahbar, S.; Pashaiasl, M. (2020). Influence of cryopreservation on structure and function of mammalian spermatozoa: An overview. *Cell Tissue Bank*, 21, (1–15).
- [25]. Fardin A, Azar P, Maryam S, Mahshad K and Saeid N. (2016). The role of antioxidants in sperm freezing: a review. *Cell Tissue Bank*, 5, (1-7).
- [26]. Gandini L, Lombardo F, Paoli D, Caponecchia L, Familiari G, Verlengia C, Dondero F, Lenzi A C. (2000). Study of apoptotic DNA fragmentation in human spermatozoa, *Hum. Reprod.* 15, 830–839.
- [27]. Garde J, Olmo A, Soler A, Espeso G, Gomendio M, Roldan E. (2007). Effect of egg yolk, cryoprotectant, and various sugars on semen cryopreservation in endangered Cuvier's gazelle (*Gazella cuvieri*). *Animal reproduction science* 108(3-4):384-401
- [28]. Gharagozloo P, Aitken R. (2011). The role of sperm oxidative stress in male infertility and the significance of oral antioxidant therapy. *Hum Reprod*, 26, 1628–1640.
- [29]. Griveau J, Dumont E, Renard P, Callegari J, Le Lannou D. (1995). Reactive oxygen species,

- lipid peroxidation and enzymatic defense systems in human spermatozoa, *J. Reprod. Fertil.* 103, 17–26.
- [30]. Henry M, Noiles, E, Gao D, Mazur P and Critser. (1993). Cryopreservation of human spermatozoa. IV. The effects of cooling rate and warming rate on the maintenance of motility, plasma membrane integrity, and mitochondrial function. *Fertil. Steril.* 60, 911–918.
- [31]. Holt W V and Abaigar T, Jabbour H (1996). Oestrous synchronization, semen preservation and artificial insemination in the Mohor gazelle (*Gazella dama mhorr*) for the establishment of a genome resource bank programme, *Reprod Fertil Dev.* 8(8):1215-22.
- [32]. Jeyendran R, Vander-Ven H, Perez-Pelaez M, Crabo B, Zanevld L. Development of an assay to assess the functional integrity of the human sperm membrane and its relationship to other semen characters. *J Repord Fertil* 1984; 70:219-228.
- [33]. Mazor D, Golan E, Philip V. (1996). Red blood cell permeability to thiol compounds following oxidative stress, *J. Eur. Haematol.* 57 241–246.
- [34]. Michael A, Alexopoulos C, Pontiki E, Hadjipavlou-Litina D, Saratsis P, Boscoc C. (2007). Effect of antioxidant supplementation on semen quality and reactive oxygen species of frozen-thawed canine spermatozoa, *Theriogenology* 68, 204–212.
- [35]. Mongioi L, Calogero A. E., Vicari E, Condorelli R. A., Russo G I, Privitera S, Morgia G, Vignera S. La. (2016). The role of carnitine in male infertility. *Andrology.* 4, (5). Pages 800-807.
- Mortimer, S. T., van der Horst, G., and [36] Mortimer, D. (2015). The future of computer-aided sperm analysis. *Asian J. Androl.* 17, 545–553
- [37]. Oeda T, Henkel R, Ohmori H, Schill WB.(1997). Scavenging effect of N-acetyl-L-cysteine against reactive oxygen species in human semen: a possible therapeutic modality for male factor infertility, *Andrologia.* 29(3):125–131.
- [38]. Perumal.P, Srivastava S., Ghosh S and BaruahK. (2014). Computer-Assisted Sperm Analysis of Freezable and Nonfreezable Mithun (*Bos frontalis*) Semen. *Journal of Animals V:* 2014, Article ID 675031, pages 1-6
- [39]. Pintado B, De La Fuente J, Roldan E. (2000). Permeability of boar and bull spermatozoa to the nucleic acid stains propidium iodide or Hoechst 33258, or to eosin: accuracy in the assessment of cell viability. *J Reprod Fertil* 118: 145-152.
- [40]. Ranjan R, Ramachandran N, Jindal S and Sinha N. (2009). Hypo osmotic swelling test in frozen thawed goat spermatozoa. *The Indian journal of animal sciences* 79(10):1022-1023
- [41]. Safarinejad M and Safarinejad S (2008). Efficacy of Selenium and/or N-Acetyl-Cysteine for Improving Semen Parameters in Infertile Men: A Double-Blind, Placebo Controlled, Randomized Study. *Journal of Urology.* 10.015
- [42]. Sariozkan S, Bucak M N, Tuncer P B, Ulutas P A, Bilgen A. (2009). The influence of cysteine and taurine on microscopic-oxidative stress parameters and fertilizing ability of bull semen following cryopreservation, *Cryobiology* 58, 134–138.
- [43]. Sariozkan S. et al (2014). Influence of various antioxidants added to TCM-199 on post-thaw bovine sperm parameters, DNA integrity and fertilizing ability. *Cryobiology* 68, 129–133.
- [44]. Sekoni VO, Kumi-Diaka J, Suror DI, Njoku CO, Olorunju SAS (1988). Seasonal and monthly variations in the incidence of morphological abnormalities in bovine spermatozoa in Shika, Zaria, Northern, Nigeria. *Anim Reprod Sci.* 17:61-67.
- [45]. Sedigheh A, Reihane B, Akram G, Azadeh N. (2016). Antioxidant supplements and semen parameters: An evidence-based review. *Int J Reprod Biomed.* 14(12):729-736.
- [46]. Sheweita S, Tilmisany A, Al-Sawaf H. (2005). Mechanisms of male infertility: role of antioxidants. *Curr Drug Metab;* 6: 495-501.
- [47]. Steven P. Brinsko D, Terry L. Blanchard D, Dickson D. Varner D, James S, Charles C. Love D. (2011). Manual of Equine Reproduction 3rd Edition.
- [48]. Storey B T. (1997). Biochemistry of the induction and prevention of lipoperoxidative damage in human spermatozoa, *Mol. Hum. Reprod.* 3, 203–213.
- [49]. Stradaoli G, Sylla L, Zelli R, Chiodi P, Monac M. (2004). Effect of L-carnitine administration on the seminal characteristics of oligoasthenospermic stallions, *Theriogenology* 62, 761–777.
- [50]. Szczesniak-Fabianczyk B, Bochenek M, Smorag Z, Ryszka F. (2003). Effect of antioxidants added to boar semen extender on the semen survival time and sperm chromatin structure, *Reprod. Biol.* 3 81–87.

- [51]. Tahereh Mardanshahi, M.Sc., 1 Nouroollah Rezaei, Ph.D., 1 , 2 Zohreh Zare, Ph.D., 1 Majid Malekzadeh Shafaroudi.(2019). Effects of L-Carnitine on the sperm parameters disorders, apoptosis of spermatogenic cells and testis histopathology in diabetic Rats. *Int J Reprod Biomed.* 17(5) 325-336.
- [52]. Varner, D. (2016) Approaches to breeding soundness examination and interpretation of results. *J. Equine Vet. Sci.*, 43 (8): S37-S44.
- [53]. Vera O, Amirat L, Diaz T, Vásquez L, Schmidt E, Desherces S, Anton M, Bencharif D, Tainturier D. (2009). Effect of semen dilution to low-sperm number per dose on motility and functionality of cryopreserved bovine spermatozoa using low-density lipoproteins (LDL) extender: Comparison to Triladyl and Bioxcell. *Theriogenology.* 1;71(6):895-900.
- [54]. Uysal O and Bucak M.N. (2007). Effect of oxidized glutathione, bovine serum albumin, cysteine and lycopene on the quality of frozen-thawed ram semen, *Acta. Vet. Brno.* 76, 383–390.
- [55]. Watson P F. (1995). Recent developments and concepts in the cryopreservation of spermatozoa and the assessment of their post-thawing function, *Reprod. Fertil. Dev.* 7, 871–891.
- [56]. Woolley D and Richardson D. (1978). Ultrastructural injury to human spermatozoa after freezing and thawing. *J Reprod Fertil*, 53(2):389-94.
- [57]. Yaniz J, Santolaria P, Marco-Aguado M, Lopez-Gatius F. (2008). Use of image analysis to assess the plasma membrane integrity of ram spermatozoa in different diluents. *Theriogenology* 70: 192-198
- [58]. Yaniz J, Mateos J, Santolaria P, (2012). Tris buffer improves fluorescence yield of ram spermatozoa when evaluating membrane integrity. *Microscop Res Techn* 75: 520-523
- Yaniz J., Palacin I, Vicente-Fiel S, Gosálvez J, Lopez-Fernandez C and Santolaria P. (2013). Evaluation of a commercial kit based on acridine orange/propidium iodide to assess the plasma membrane integrity of ram sperm. *Spanish Journal of Agricultural Research* ,11(2), 362-365.
- [60]. Zafarullah M, W. Li Q , Sylvester J & Ahmad M . (2003). Molecular mechanisms of N-acetylcysteine actions. *Cellular and Molecular Life Sciences.* volume 60, pages6–20.